

VIRTUAL COLONOSCOPY

HÔPITAL BROME-MISSISQUOI-PERKINS



Your physician has prescribed a virtual colonoscopy examination. This computed tomography (CT) scan produces 3D images of the internal lining of the large intestine (colon and rectum) for the identification of colorectal pathologies.

ABOUT THE EXAMINATION

- These examinations are by appointment only.
- No sedative or anesthetic is required for this procedure.
- The examination lasts approximately 45 minutes.

ALLERGIES | DIABETES

If you are diabetic or allergic to iodinated contrast agents (iodine):

- Inform your doctor. He will give you specific instructions to follow.
- Notify the administrative officer of the medical imaging department at the number in the checklist below.
- Upon arriving at the medical imaging department, notify the technologists of your diabetes and/or allergies.

PREPARING FOR THE PROCEDURE

Preparing for the procedure is complex but essential to the quality of the scan.

Preparation instructions must be followed to the letter to eliminate fecal residues that might be confounded with other pathologies. Do not make any changes or substitutions without your doctor's authorization.

THE PREPARATION INSTRUCTIONS ARE ON THE NEXT PAGE.

ON THE DAY OF THE EXAMINATION

- Plan to spend 45 to 60 minutes for the duration of your examination.
- You will be asked to put on a hospital gown.
- The technologist might inject an antispasmodic (e.g., Buscopan).
- A small plastic tube will be inserted into your anus so that carbon dioxide (CO₂) can be pumped into your colon. This will inflate your colon, making it easier to view the lining. This action may cause discomfort (bloating sensation). You will then be asked to perform certain movements.
- A CT scan will then be conducted. The radiologist might request an iodinated contrast injection.
- Do not hesitate to ask the technologists any question you might have during the examination.

BRING THE FOLLOWING TO THE EXAMINATION

- A valid health insurance card (to avoid paying for this test).
- A valid hospital card (if needed, you will have to go to the reception and admitting department to get a card before your appointment).
- A list of your medications, if requested.
- A bag for your personal items. Do not bring or wear any jewels or valuables. (The hospital is not responsible for lost or stolen objects.)

CANCELLING OR POSTPONING AN APPOINTMENT

To cancel or postpone your appointment, refer to the coordinates in the memory aid below.

DO YOU HAVE ANY QUESTIONS?



santeestrie.qc.ca/imaging



Info-Santé 811 OR
Speak to your prescriber.

MEMORY AID

My examination date: _____

Hôpital Brome-Missisquoi-Perkins

950, rue Principale, Cowansville (Québec) J2K 1K3
Radiology reception desk situated on the 1st floor.

To cancel or postpone your examination:
450-266-4342, Ext. 65800

PROCEDURES TO FOLLOW FOR THE VIRTUAL COLONOSCOPY

→IF YOU ARE FOLLOWING A SPECIFIC DIET, CONFIRM WITH YOUR DOCTOR THAT THIS PREPARATION IS SUITABLE.←

LIST OF REQUIRED PRODUCTS:

- One Pico-Salax box containing 2 powder sachets: Purchase this product at your pharmacy (\$).
- One bottle of E-Z-CAT 225 ml (non-iodinated contrast medium): The radiology department will supply this product.
- One bottle of Gastrographin 30 ml (iodinated contrast medium): The radiology department will supply this product.
Inform us if you are allergic to this product.

PREPARATION AND DIET TO BE FOLLOWED:

48 HOURS BEFORE THE PROCEDURE:

Date: _____

- **Unrestricted low-residue diet** (see *LOW-RESIDUE DIET* on next page).
- All liquid diet foods are also permitted (see *LIQUID DIET* on next page, *unrestricted foods*).
- Drink one glass of water or clear liquid every hour until bedtime.

24 HOURS BEFORE THE PROCEDURE:

Date: _____

- **Unrestricted liquid diet** for all meals and snacks (see *LIQUID DIET* on next page).
- Drink one glass of water or clear liquid every hour until bedtime.
- During preparation, if nausea occurs, use Graval® as needed.

TAKING THE VARIOUS PRODUCTS AT THE PRESCRIBED TIME

TIME	INSTRUCTIONS
12:00 P.M. (noon)	<ul style="list-style-type: none">• Dissolve the first sachet of Pico-Salax in 150 ml of cold water, mix for 2-3 minutes, and drink.• Drink 250 ml (8 oz) of water every hour for 4 to 6 hours after taking a sachet of Pico-Salax.
7:00 P.M.	<ul style="list-style-type: none">• Dissolve the second sachet of Pico-Salax in 225 ml of E-Z-Cat. Stir and drink it all.• Drink 250 ml (8 oz) of water every hour for 4 to 6 hours after taking a sachet of Pico-Salax.
9:00 P.M.	<ul style="list-style-type: none">• Mix 1 bottle (30 ml) of Gastrographin diluted in 250 ml of water.• Drink this mixture.
12:00 A.M. (midnight)	<ul style="list-style-type: none">• Do not drink or eat until the colonoscopy is completed.

ON THE DAY OF THE PROCEDURE

- Be fasting since midnight.
- Do not drink, eat, or smoke.
- For taking medications only: A sip of water is permitted.

LOW-RESIDUE DIET	LIQUID DIET
<ul style="list-style-type: none"> ✓ White bread without fibre and without butter (fibre: check the nutritional value table on the package) ✓ Hardboiled eggs or poached eggs with firm yolks ✓ Cooked fish without added fat (canned tuna in water is permitted) ✓ Peeled boiled potatoes ✓ White rice ✓ White pasta (can be cooked in chicken broth) ✓ Salt and pepper 	<ul style="list-style-type: none"> ✓ Water ✓ Gatorade or an equivalent clear-coloured beverage (EXCEPT red). These beverages are particularly recommended as they maintain hydration and improve bowel preparation. ✓ Pulp-free fruit juice (apple juice, orange juice, white grape juice, or lemonade) ✓ Clear soft drinks (Seven-Up, Sprite) ✓ Clear broths or consommés ✓ Ensure or fibre-free equivalent beverage (vanilla flavour only) ✓ Jell-O (except red and purple) ✓ Popsicle ✓ Tea, herbal tea and coffee (without cream or milk) in limited amounts. Sugar and honey are permitted.
AVOID	
<ul style="list-style-type: none"> ✗ Dairy products (yogurt, cheese, etc.) ✗ Fruits and vegetables ✗ Butter and other fats ✗ Nuts ✗ Jam ✗ Solid meat 	<ul style="list-style-type: none"> ✗ All red liquids (e.g., cranberry juice, tomato juice, red Gatorade, red and purple Jell-O).



REMINDER FOR DIABETICS:

Consult your doctor for instructions to adjust your medication.

If you are taking medications,
you can have a sip of water as usual to swallow them.

Author

Équipe de l'imagerie médicale et de médecine nucléaire

Revision and layout

Service des communications et des relations médias
Présidence – direction générale

© Centre intégré universitaire de santé et de services sociaux de l'Estrie –
Centre hospitalier universitaire de Sherbrooke

santeestrie.qc.ca

January 2025 – 4-6-12396 (French version) | 4-6-12397 (English version)