

BI-PEGLYTE®

SEVEN DAYS BEFORE THE COLONOSCOPY

Seven days before your exam, stop taking your iron supplements.

THREE DAYS BEFORE THE COLONOSCOPY

For three days before your examination, avoid the following foods:

- Cold cuts and sausages
- Raw vegetables
- These cooked vegetables: corn, cabbage family (broccoli, cauliflower, Brussels sprouts, cabbage, kale, chard, etc.)
- Fruit with small pits or seeds: strawberries. raspberries, kiwis, watermelon, blackberries, grapes, tomatoes, etc.
- THE DAY BEFORE THE COLONOSCOPY

- Whole grains, nuts, sesame and flax seeds
- Legumes (dried beans, lentils, peas, etc.)
- Whole wheat or multigrain foods (multigrain • breads, brown or wild rice, whole wheat pasta, quinoa, etc.)

The day before your examination, do not eat any food. You can't eat, but you can drink.

As soon as you get up, consume only clear liquids throughout the day. Drink plenty desired liquids allowed.

ALLOWED	PROHIBITED
Water and ice;	No red, purple, blue, or green coloured drinks;
 Apple, white grape or clear white cranberry juice without pulp; 	No alcoholic beverages;
 Jell-0 (except red and purple); 	No almond or soy milk;
 Beef or chicken consommé or broth (without noodles, vegetables, or solid residue); 	No opaque beverages;
 Herbal tea, coffee, and tea (without cream or milk); 	 No dark soft drinks (Coke, Pepsi);
Gatorade-type energy drinks (except red and purple);	No dairy products or alternatives (Ensure, Boost)
Clear soft drinks (7-Up, Sprite, ginger ale).	No liquid with pulp.

Do not eat any food and do not drink milk the day before and the day of the colonoscopy.







DATE: __/ __/ __

THE DAY BEFORE THE COLONOSCOPY



THE MORNING

Prepare the Bi-PegLyte® solution as follows:

- 1. Pour the contents of 1 full pouch into 1 litre of water (32 ounces).
- 2. Shake for 2 to 3 minutes until the solution becomes clear. Set it aside.
- 3. In another container, pour 1 full pouch into 1 litre of water.
- 4. Shake for 2 to 3 minutes until the solution becomes clear.
- 5. Refrigerate **both solutions** as that improves the taste.

THE AFTERNOON

6. Around 3:00 p.m., take the 3 bisacodyl tablets with water. Do not chew or crush the tablets.



Do not take any antacids within one hour of taking the tablets.



THE EVENING

7. Around 7:00 p.m., quickly drink 1 glass of the first solution (about 250 mL) every 10 minutes, until you have drunk the entire litre of solution (total of 4 glasses of 250 mL).

Check off each glassful as you finish it.



4

You should begin to have watery bowel movements within an hour and continue having them for 1 to 2 hours after drinking all of the solution.

2

THE DAY OF THE COLONOSCOPY



Do not eat any food. You can drink clear liquids allowed for up to 4 hours before your appointment.



8. Four hours before your appointment, quickly drink 1 glass of the second solution (about 250 mL) every 10 minutes until you have drunk the entire litre of solution (total of 4 glasses of 250 mL). 1 2 3

Check off each glassful as you finish it.

Arrange to have someone take you home after your colonoscopy.

Precautions

- Bloating or abdominal distension may occur before the first stool is passed. If abdominal distension or discomfort persists, temporarily stop drinking the solution or take longer to drink each 250 mL serving (more than 10 minutes) until the symptoms pass.
- If you experience severe abdominal swelling, pain, or distension, temporarily slow down or stop taking the solution until the symptoms pass. Report these symptoms to your doctor.
- Tell your doctor immediately if you have severe abdominal pain or bleeding from the rectum.

Authors

Gastroenterology and endoscopy teams

Updated Dr. Julie Carrier and Dr. Nathalie McFadden Marie-Pier Lescault, Clinical Nurse Specialist Annie Paguin, Head, Endoscopy Department

Revision and Layout

Service des communications

Direction des ressources humaines, des communications et des affaires juridiques

© Centre intégré universitaire de santé et de services sociaux de l'Estrie - Centre hospitalier universitaire de Sherbrooke, 2020

santeestrie.qc.ca

December 2022 - English version : 1-6-71747- French version : 1-6-71728

Centre intégré universitaire de santé et de services sociaux de l'Estrie – Centre hospitalier universitaire de Sherbrooke

